

METHOD FOR PREPARING ZINC-OLIGOPEPTIDE EASILY ABSORBABLE BY THE HUMAN BODY

BACKGROUND OF THE INVENTION

5

The present invention relates to a method for preparing zinc-oligopeptides which can be easily absorbed by the human body. Also, the present invention is concerned with a beverage or food composition, which can easily
10 provide zinc for the human body.

Loss of the mineral balance in the body has been found as one of the causes of adult diseases, which have recently been increasing in incidence. With growing older, the human body becomes poorer at absorbing minerals. Nowadays, a
15 deluge of processed foods is becoming as one main cause of upsetting the mineral balance of the body because most of them contain materials inhibiting mineral absorption.

Of various minerals, zinc is involved particularly in the onset of diabetes mellitus, one of the most popular
20 diseases today. In fact, not only adults, but also a surprising number of children suffer from the disease at present, which is believed to be attributed in part to the lack of zinc. In Oriental medicine, zinc is also described to have physiological activity associated with sugar control
25 and vigor in the body. Deeply affected in diabetes mellitus

patients, the blood sugar control of the body has direct influence on the energy production necessary for life. Although other physiological functions are active, abnormal regulation of blood sugar in the body, which means that the metabolic process of converting sugar into energy is in an abnormal state, lowers the immunity of the body to exogenous pathogens as well as latent viruses, resulting in the body falling ill.

In the body, zinc serves as an essential mineral in activating insulin, in addition to being involved in regenerating muscle tissues and nerve tissues. Abundant as it is in blood, insulin cannot exert its full effects in the absence of zinc. That is, insulin which is not associated with zinc is not beneficial to diabetics. A similar case can be found with amylase, which is unable to function as a saccharification catalyst unless it is associated with ionic calcium.

One of the most important physiological activities of zinc is to activate insulin into a form useful in the treatment of diabetes mellitus. In turn, the activated insulin is responsible, in great part, for the production of energy. In addition, zinc was found to inhibit the expression of mutant genes, thereby making a contribution to anticancer activity. Further, zinc is known to take part in a catalytic reaction necessary for DNA polymerization and

therefore affect the rapid regeneration of injured tissue. In this regard, zinc has some connection with acceleration of wound healing, prevention of prostate problems and hair loss, and treatment of acne and rheumatoid arthritis.

5 Recently, sufficient intake of zinc has been reported to significantly decrease the incidence of disease in children. When their diets are supplemented with zinc, children are 40 % less likely to be taken ill with pneumonia and 25 % less likely to get diarrhea. As stated above,
10 diabetes mellitus may be caused when dietary zinc is insufficient. When insulin is not activated, the body is significantly deprived of available sugar, leading to loss of vigor.

 In order to be activated in association with zinc, the
15 insulin must not be in a pro-insulin form, but in a functional form. Stoichiometrically, one molecule of functional insulin (molecular weight 6,615) associates with a zinc atom (atomic weight 65). Therefore, functional insulin must be associated with zinc at a weight ratio of
20 approximately 1,000:1. In the medical world, globulin zinc I is used as an insulin formulation, which is generally for subcutaneous injection, for the purpose of zinc activation of insulin.

 Occurrence of diabetes mellitus in children in recent
25 times, which was rare in the past, is believed to be

strongly affected by dietary lifestyles, but not heredity. Fried foods, which children usually like, contain a large quantity of fat that suppresses the activity of zinc. Also, a high intake of processed foods inhibits activities of calcium as well as zinc because of their high contents of phosphoric acid. Lipids and phosphates are known to actively inhibit the absorption of minerals irrespective of which form they have.

Naturally, living bodies have to supplement consumed or deficient materials by themselves. Hence, it is necessary that people exercise restraint in their ingestion of materials which inhibit such a natural supplementary function. The loss of the supplementary functions owing to ingestion of inhibitory materials causes a vicious cycle of deficiency. For instance, deficient dietary zinc lowers the activity of insulin, and unconsumed insulin causes a decrease or ceasing of the production of insulin in the pancreas. On the other hand, when insulin is actively consumed, it must be replaced, and thus insulin is more actively produced in the pancreas. This is demonstrated by the fact that athletes, who consume large energy, contain zinc at an amount 20 times as much as that of ordinary people. Zinc enables insulin to actively promote the metabolism of sugar in the body.

However, most of the mineral-enriched materials

developed thus far, are poor in absorbability by the human body. Calmodulin, a calcium-chelated oligopeptide, was found in neurotic system of the human body, and to be absorbed easy as Calmodlin molecular structure by the small intestine.

SUMMARY OF THE INVENTION

Hinted by the structure of calmodulin, the intensive and thorough research on facilitating of the human body to absorb zinc, conducted by the present inventors, resulted in the finding that minerals are better absorbed by the body when they combine with organic materials, especially oligopeptides, rather than alone.

Therefore, it is an object of the present invention to provide a method for preparing a zinc-oligopeptide which can be easily absorbed by the body.

It is another object of the present invention to provide use of zinc-oligopeptide in foods.

In accordance with an aspect of the present invention, there is provided a method for preparing a zinc-oligopeptide easily absorbable by the body, comprising the steps of: proteolyzing a suspension of protein in deionized water at a neutral pH range in the presence of a protease to give a mixture of oligopeptides; chelating zinc ions with the

oligopeptides to give a zinc-oligopeptide solution; concentrating the zinc-oligopeptide solution and drying the concentrate to a powder.

In accordance with another aspect of the present invention, there is provided a beverage comprising the zinc-oligopeptide, in combination with at least one ingredient selected from the group consisting of vitamin -C, vitamin-B₁, vitamin-B₂, fructose, α -amylase decomposed starch and magnesium stearate.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows a structure of an oligopeptide associated with a zinc ion.

Fig. 2 shows a structure of an oligopeptide alone.

DETAILED DESCRIPTION OF THE INVENTION

In accordance with one embodiment of the present invention, there is provided a method for preparing zinc-oligopeptide. First, a suspension of protein in deionized water is subjected to proteolysis in a neutral pH range for a long period of time to give oligopeptides. These are used to chelate zinc ions to form zinc-oligopeptides. The resulting solution is concentrated and dried to yield zinc-

oligopeptide powder.

In detail, 100 weight parts of protein are suspended in 800 weight parts of deionized water and added with 2-4 weight parts of protease. Proteolytic reaction is conducted at pH 3.5-6.0 for 10-12 hours to give oligopeptides. Based on 1,000 weight parts of the oligopeptide thus obtained, one weight part of zinc ions is mixed and allowed to chelate, to yield a zinc-oligopeptide. Then, the resulting zinc-oligopeptide solution is concentrated to a solid content of 32-36% and dried to produce zinc-oligopeptide powder.

Either vegetable protein or animal protein may be used for obtaining oligopeptides.

In accordance with another embodiment of the present invention, there is provided a zinc-oligopeptide-containing beverage. To this end, vitamin-C, vitamin-B₁, vitamin-B₂, fructose, α -amylase decomposed starch, and/or magnesium stearate may be mixed with liquid zinc-oligopeptide. Alternatively, this composition is dehydrated to give powder suitable for use in capsules or tablets.

More specifically, 99.5 % of the zinc-oligopeptide is mixed with 0.01-0.05 % of vitamin-C, 0.01-0.05 % of vitamin-B₁, 0.01-0.05 % of vitamin-B₂, 4.0-5.0 % of α -amylase decomposed starch, and/or 0.01-0.05 % of magnesium stearate.

A better understanding of the present invention may be obtained in light of the following examples which are set

forth to illustrate, but are not to be construed to limit the present invention.

EXAMPLE 1

5

As an oligopeptide source, a vegetable protein such as bean protein and gluten, or an animal protein such as casein and gelatin was used. 100 weight parts of protein with a purity of 95 % or higher was suspended in 800 weight parts
10 of water and allowed to undergo proteolysis in the presence of 3 weight parts of protease at pH 3.5-6.0 for 12 hours to give oligopeptides. Of the proteolysates, non-water soluble ones were filtered off. One weight part of zinc ions was added based on 1,000 weight parts of the oligopeptide to
15 form a zinc-oligopeptide composite. Then, the resulting solution was concentrated to a solid content of 35 % and spray-dried to give a zinc-oligopeptide powder.

EXAMPLE 2

20

The zinc-oligopeptide solution prepared in Example 1 was controlled to have a concentration of 28 % and formulated as indicated in Table 1, below, to provide a zinc-oligopeptide beverage with a full recommended dietary
25 allowance (RDA) of zinc (12-15 mg/day).

TABLE 1
Composition of Zinc-Oligopeptide Beverage

Ingredients	Amounts (wt%)
28% Zn-Oligopeptide Sol'n	1.0
Vitamin-B ₁	0.01
Fructose	12
Vitamin-C	0.03
Vitamin-B ₁₂	0.01
Water	86.95
Total	100

5

EXAMPLE 3

The zinc-oligopeptide prepared in Example 1 was formulated with other ingredients as shown in Table 2, below, and 0.5 g of the resulting mixture was filled into a capsule.

TABLE 2
Composition of Zinc-Enriched Capsule

Ingredients	Amounts (wt%)
Zn-Oligopeptide Powder	99.95
Vitamin-B ₁	0.01
Vitamin-C	0.03
Vitamin-B ₁₂	0.01
Total	100

15

EXAMPLE 4

The zinc-oligopeptide prepared in Example 1 was formulated with other ingredients as shown in Table 3, below, and 0.5 g of the resulting mixture was formed into a zinc-enriched tablet.

TABLE 3
Composition of Zn-Enriched Tablet

Ingredients	Amounts (wt%)
Zn-Oligopeptide Powder	94.95
Vitamin-B ₁₂	0.01
α -Amylase Decomposed starch	4.7
Vitamin-C	0.03
Vitamin-B ₆	0.01
Mg Stearate	0.3
Total	100

10

As described hereinbefore, the zinc-oligopeptide of the present invention has a molecular weight of from 800 to 1,000. Because its molecular weight is smaller than the average molecular weight (24,000-28,000 of membrane integral proteins of the small intestine, through which molecules pass in and out of the cell, the zinc oligopeptide of the present invention can be readily absorbed by the body. Additionally, zinc is chelated by water-soluble oligopeptides, so that its absorption by the body is not

inhibited by other compounds present in the digestive tract.

The present invention has been described in an illustrative manner, and it is to be understood that the terminology used is intended to be in the nature of
5 description rather than of limitation. Many modifications and variations of the present invention are possible in light of the above teachings. Therefore, it is to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically
10 described.